

Figure 1/21

Construction of intracellular signalling chimera:

1. PCR myristoylated CD3ζ





2. Cut and clone PCR fragment

*The MZE series contains a 9aa HA epitope at the 3' end.







From plasmid #FK12/KS

3. SEQUENCE insert

4. Cut at XhoI or SalI and add FKBP domains



plasmid #MZFn



plasmid #MFnZ

Figure 2/21

Construction of extracellular signaling chimera:

1. PCR murine signal peptide

2. PCR CD3 trans-membrane and cytoplasmic domains



Figure 3A/21

3. PCR FKBP12



Figure 3B/21



homology

SacII

6048: 5'-CGACACCGCGGCCACCATGGCCACAATTGGAGC-3'

Kozak M A T I G

homology

XhoI

6049: 5'-CGACACTCGAGAGCCCATGACTTCTGG

L A W S _ ?





Asp-Gyl#1

homology

XhoI *

6050: 5'CGACACTCGAGCTCTGCTACTTGCTAGGTGGAATCCTCTTC-3'

E L C Y L L G G I L F

*A to G

homology

EcoRI *

6051: 5' GCGAATTCTTAGCGAGGGGCCAGC-3'

St R P A L *G to C

3'Sal #B

homology

EcoRI SalI

8462: 5' GCGAATTCTTAGTCGACGCGAGGGGCCAGGGTC-3'

St R PAL

Cys-Gly #2

homology

XhoI *

7129: 5' GGGCTCGAGCTCGGCTACTTGCTAG-3'

L G Y L L *T to G



Figure 4B/21


homology
Asp-Lys #4
XhoI
* *
8061: 5'-CGACACTCGAGCTCTGCTACTTGCTAAAGGGAATCCTCTTC-3'
E L C Y L L K G I L F
*GATtoAAG
homology
#4 Extension
XhoI
8907: 5'-CGACACTCGAGCTGCTGGATCCGAAGCTCTGCTACTTGCTAAAG-3'
E L L D P K L C Y L L K
homology
TAC-Tm #3
XhoI
7220: 5'-CGACACTCGAGACAACAGAGTACCAGGTAGC-3'
E T T E Y Q V
immunophilin
FKBP12
homology
XhoI
6052: 5'-CGACACTCGAGGGCGTGCAGGTGGAGAC-3'
E G V Q V E
homology
SalI
6053: 5'-CGACAGTCGACTTCCAGTTTTAGAAGC-3'
V E L K L L
FKBP13
homology
XhoI
8460: 5'-TCGACACTCGAGACGGGGGCCGAGGGC-3'
E T G A E G
homology
SalI
8461: 5'-CCGACAGTCGACCTCTATTTTGAGCAGC-3'
V E I

Figure 4C/21



Figure 5/21



Figure 6A/21



MZF3E



MZE

Cut XhoI/SalI; CIP; + FKBP12 X 3



MF3E







Figure 6B/21

Activity of FK1012A on the chimeric FKBPX3/CD3 zeta receptor



Figure 7/21

Activation of an NFAT reporter via signalling through a myristoylated CD3 zeta/FKBP12 chimera



Figure 8/21



Figure 9A (#1)/21



13



12

Figure 9A(#2)/21



Figure 9B (#1)/21



16



17

FK506

Figure 9B (#2)/21

Scheme 1



Figure 10/21

Scheme 2: Synthesis of Dimers



TBS-OTf



1) 2M NH3 / MeOH

2) formic acid / acetic anhydride or acetic anhydride



SeO_2

Lit ref: N.H. Sigal & F.J. Dumont, Ann. Rev. Immunol. p519 1992

Lit refs: D.K. Donald et.al. Tetrahedron Letters p1375, 1991, P.Kocovsky, Tetrahedron Letters p5521, 1992



N,N'-Disuccinimidyl carbonate, Et_3N, CH_3CN

Figure 11A/21





Figure 11B (#1)/21

An additional modified FK520 (FK1040) that interferes with FKBP12 yet should bind the FKBP12 mutant: F36A or F99A or Y26A, or combinations thereof is



Figure 11B (#2)/21

Scheme 3 Heterodimerization



Figure 12/21

Scheme 3: Synthesis of heterodimers



In this example, a heterodimer of a cyclosporine analog and FK520A-NHCO-R were heterodimerized. However, the scheme can easily incorporate other FK506/520 derivatives to form hetero or homodimers

Figure 13/21



Figure 14/21


Cytoplasm
Myristate
FKBP12
FK1012
FKBP12
FK1012
Tyrosine Kinases
PLCγ1
Ca2+
DAG
Calcineurin
PKC, MAPK
NFATc
NFATn
NFAT
NFAT
NFAT
Inducible gene

Figure 15/21

A FK1012s



B FK506 Monomer with a C10 Bump



Figure 16 (#1)/21

C FK506 Monomer with a C9 Bump



D HED Reagent Synthesis



Figure 16 (#2)/21


1 Ac2O
2 NBS, AIBN
3 NEt4OAc
4 NaOMe
1 2 equiv
(Im)2CO
2
NH2
H2N
cyclosporin A (CsA)
(CsA)2

Figure 17/21

A cDNA construct



Figure 18A/21

B expressed protein



Figure 18B/21



Figure 19/21



Figure 20A/21



Figure 20B/21


FK506
1 TBSOTf
2 DIBAL-H
1 PhCH2N2
2 HF, CH3CN
R=CH2Ph
1 Ph3Bi(OAc)2
2 HF, CH3CN
R=Ph
1 TBSOTf
2 NaBH4
3 DIBAL-H
PhCHN2
1 TBSOTf
2 CH2N2
3 TFAOH
1 CO(Im)2
2 HF, CH3CN
HF
CH3CN
R = H, Me
n = 2,4,6,8
Target
HED
Reagent

Figure 21/21